


How to test your drugs for fentanyl

with BTNX Rapid Response™ test strips

It is most accurate to test everything you intend to consume.

After testing, you can either drink the mixture, snort it, or wait for the water to evaporate to use. You can also test a sample, but fentanyl isn't always mixed evenly into drugs.

1 Prepare drugs



Powder, Crystal, Pills

Crush up and mix well. Measure out 10 mg of powder into a small baggie or container like a water bottle cap.

10 mg ≈ the size of a grain of rice or 1 micro scoop



Injection

Prepare shot in an unused cooker; set aside. Use cooker with residue.

Whole supply test

Calculate entire supply weight. Crush up & mix well.

2 Add water

Add about 5 ml/cc of water to baggie, container, or cooker. Mix well.


5 ml/cc water ≈ 1 plastic bottle cap or teaspoon



Whole supply test

Drug weight	Add water & mix well	
	MOST DRUGS	METH OR MDMA
10 mg	5 ml/cc or 1 tsp	¼ cup or 2 oz
.1 g	50 ml/cc or 10 tsp	2.5 cups or 20 oz
.5 g	250 ml/cc or 1 cup	12.5 cups or 100 oz

3 Dip strip for 15 seconds



Hold strip's dark blue end. Dip other end into mixture for 15 seconds.

Don't dip past the wavy blue lines.

4 Wait 5 minutes

Lay the test strip down on a flat surface.

Wait 5 minutes to read result.



5 Read result



1 red line = POSITIVE for fentanyl
Fentanyl may be present in the drugs.

If there is only a red line on the left or no red lines appear, the test is invalid. Redo with a new strip.



2 red lines = NEGATIVE for fentanyl*
Look closely for a second line; it can be faint!

*Just because a portion of your supply tested negative does not mean your entire supply is fentanyl free.

Take steps to lower your risk

Even if a fentanyl test result is negative, there is still a risk of an overdose. You can take these steps to lower that risk.

Test your drugs.
Never use alone.
Carry naloxone.



Always have naloxone visible when using

Use slow, use less

Overdoses can happen quickly. Space out doses.

Consider alternatives to injecting

Snorting or smoking carries a reduced overdose risk compared to injecting.

Take turns when you use drugs

Make sure one person is alert and naloxone is visible.

Test your drugs every time for fentanyl

Always test your drugs even if you got them from the same person or same bag.

Reach out if you use alone

Call a friend or use a service like **Never Use Alone** at 877-696-1996 or **The Brave App**.

What is fentanyl?

Fentanyl is a strong synthetic opioid that increases the risk of an overdose.

Powdered fentanyl looks just like many other drugs. It may be mixed with drugs like heroin, cocaine, methamphetamine, ketamine, MDMA, or in fake pills that look like prescription pills.

Save Lives
OREGON

SaveLivesOregon.org

What should I know about fentanyl test strips?

Fentanyl test strips can be helpful, but they are NOT perfect tools.

- Depending on the drugs you are testing and/or the method used to test, your results may not be correct or reliable.
- False positives and false negatives can occur.

- Fentanyl test strips can tell you if your drugs contain fentanyl but NOT how much OR how strong the effect of the drugs will be on you.
- Just because a portion of your supply tested negative does not mean your entire supply is fentanyl free.